

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 401/14, A61K 31/44	A1	(11) International Publication Number: WO 97/17337 (43) International Publication Date: 15 May 1997 (15.05.97)
(21) International Application Number: PCT/GB96/02684 (22) International Filing Date: 4 November 1996 (04.11.96) (30) Priority Data: 9523064.5 10 November 1995 (10.11.95) GB (71) Applicant (for all designated States except US): MERCK SHARP & DOHME LIMITED [GB/GB]; Hertford Road, Hoddesdon, Hertfordshire EN11 9BU (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): BOURRAIN, Sylvie [FR/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). MACLEOD, Angus, Murray [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). NEDUVELIL, Joseph, George [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). SHOWELL, Graham, Andrew [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). (74) Agent: THOMPSON, John; Merck & Co., Inc., European Patent Dept., Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: SUBSTITUTED 1-INDOLYLPROPYL-4-BENZYL-TETRAHYDROPYRIDINE DERIVATIVES (57) Abstract A class of 1-[3-(1 <i>H</i> -indol-3-yl)propyl]-4-benzyl-1,2,5,6-tetrahydropyridine derivatives, substituted at the 5-position of the indole nucleus by a 1,2,4-triazol-4-yl moiety, and on the methylene linkage of the benzyl moiety by an alkyl, alkoxy or alkoxy-alkoxy substituent, are selective agonists of 5-HT ₁ -like receptors, being potent agonists of the human 5-HT _{1D} receptor subtype whilst possessing at least a 10-fold selective affinity for the 5-HT _{1D} receptor subtype relative to the 5-HT _{1D} subtype; they are therefore useful in the treatment and/or prevention of clinical conditions, in particular migraine and associated disorders, for which a subtype-selective agonist of 5-HT _{1D} receptors is indicated, whilst eliciting fewer side-effects, notably adverse cardiovascular events, than those associated with non-subtype-selective 5-HT _{1D} receptor agonists.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

SUBSTITUTED 1-INDOLYLPROPYL-4-BENZYL-
TETRAHYDROPYRIDINE DERIVATIVES

The present invention relates to a class of substituted
5 tetrahydropyridine derivatives which act on 5-hydroxytryptamine (5-HT)
receptors, being selective agonists of so-called "5-HT₁-like" receptors. They
are therefore useful in the treatment of clinical conditions for which a
selective agonist of these receptors is indicated.

It has been known for some time that 5-HT₁-like receptor agonists
10 which exhibit selective vasoconstrictor activity are of use in the treatment
of migraine (see, for example, A. Doenicke *et al.*, *The Lancet*, 1988, Vol. 1,
1309-11; and W. Feniuk and P.P.A. Humphrey, *Drug Development*
Research, 1992, 26, 235-240).

The human 5-HT₁-like or 5-HT_{1D} receptor has recently been shown
15 by molecular cloning techniques to exist in two distinct subtypes. These
subtypes have been termed 5-HT_{1D α} (or 5-HT_{1D.1}) and 5-HT_{1D β} (or
5-HT_{1D.2}), and their amino acid sequences are disclosed and claimed in
WO-A-91/17174.

The 5-HT_{1D α} receptor subtype in humans is believed to reside on
20 sensory terminals in the dura mater. Stimulation of the 5-HT_{1D α} subtype
inhibits the release of inflammatory neuropeptides which are thought to
contribute to the headache pain of migraine. The human 5-HT_{1D β} receptor
subtype, meanwhile, is located predominantly on the blood vessels and in
the brain, and hence may play a part in mediating constriction of cerebral
25 and coronary arteries, as well as CNS effects.

Administration of the prototypical 5-HT_{1D} agonist sumatriptan
(GR43175) to humans is known to give rise at therapeutic doses to certain
adverse cardiovascular events (see, for example, F. Willett *et al.*, *Br. Med.*
J., 1992, 304, 1415; J.P. Ottervanger *et al.*, *The Lancet*, 1993, 341, 861-2;
30 and D.N. Bateman, *The Lancet*, 1993, 341, 221-4). Since sumatriptan
barely discriminates between the human 5-HT_{1D α} and 5-HT_{1D β} receptor

subtypes (cf. WO-A-91/17174, Table 1), and since it is the blood vessels with which the 5-HT_{1D β} subtype is most closely associated, it is believed that the cardiovascular side-effects observed with sumatriptan can be attributed to stimulation of the 5-HT_{1D β} receptor subtype. It is accordingly

5 considered (cf. G.W. Rebeck *et al.*, *Proc. Natl. Acad. Sci. USA*, 1994, 91, 3666-9) that compounds which can interact selectively with the 5-HT_{1D α} receptor subtype, whilst having a less pronounced action at the 5-HT_{1D β} subtype, might be free from, or at any rate less prone to, the undesirable cardiovascular and other side-effects associated with non-subtype-selective

10 5-HT_{1D} receptor agonists, whilst at the same time maintaining a beneficial level of anti-migraine activity.

The compounds of the present invention, being selective 5-HT₁-like receptor agonists, are accordingly of benefit in the treatment of migraine and associated conditions, e.g. cluster headache, chronic paroxysmal

15 hemicrania, headache associated with vascular disorders, tension headache and paediatric migraine. In particular, the compounds according to this invention are potent agonists of the human 5-HT_{1D α} receptor subtype. Moreover, the compounds in accordance with this invention have been found to possess at least a 10-fold selective affinity for

20 the 5-HT_{1D α} receptor subtype relative to the 5-HT_{1D β} subtype, and they can therefore be expected to manifest fewer side-effects than those associated with non-subtype-selective 5-HT_{1D} receptor agonists.

Several distinct classes of substituted five-membered heteroaromatic compounds are described in published European patent

25 application 0497512, and published International patent applications 93/18029, 94/02477 and 94/03446. The compounds described therein are stated to be agonists of 5-HT₁-like receptors, and accordingly to be of particular use in the treatment of migraine and associated conditions. None of these publications, however, discloses nor even suggests the


30 substituted tetrahydropyridine derivatives provided by the present invention.

Moreover, nowhere in the prior art mentioned above is there any disclosure of a subtype-selective 5-HT_{1D} receptor agonist having a 5-HT_{1D α} receptor binding affinity (IC₅₀) below 50 nM and at least a 10-fold selective affinity for the 5-HT_{1D α} receptor subtype relative to the 5-HT_{1D β} subtype.

5 The compounds according to the present invention are subtype-selective 5-HT_{1D} receptor agonists having a human 5-HT_{1D α} receptor binding affinity (IC₅₀) below 50 nM, typically below 10 nM and preferably below 1 nM; and at least a 10-fold selective affinity, typically at least a 50-fold selective affinity and preferably at least a 100-fold selective affinity,
10 for the human 5-HT_{1D α} receptor subtype relative to the 5-HT_{1D β} subtype. Moreover, the compounds in accordance with this invention possess interesting properties in terms of their efficacy and/or bioavailability.

The present invention provides a compound of formula I, or a salt or prodrug thereof:


15



(I)

wherein

20 R¹ represents hydrogen, halogen, trifluoromethyl, C₁₋₆ alkoxy or a group of formula (a):



(a)

R² and R³ independently represent hydrogen, halogen, trifluoromethyl or C₁₋₆ alkoxy; and

R⁴ represents C₁₋₆ alkyl, C₁₋₆ alkoxy or C₁₋₆ alkoxy(C₁₋₆)alkoxy.

The compounds in accordance with the present invention are
5 encompassed within the generic scope of co-pending International Patent Application No. PCT/GB95/01129, published as WO 95/32196 on 30 November 1995. There is, however, no specific disclosure therein of compounds corresponding to those of formula I above wherein R¹, R², R³ and R⁴ are as defined above.

10 As used herein, the expression "C₁₋₆ alkyl" includes methyl and ethyl groups, and straight-chained or branched propyl, butyl, pentyl and hexyl groups. Particular alkyl groups are methyl, ethyl, *n*-propyl, isopropyl and *tert*-butyl. Derived expressions such as "C₁₋₆ alkoxy" are to be construed accordingly.

15 The term "halogen" as used herein includes fluorine, chlorine, bromine and iodine, especially fluorine.

For use in medicine, the salts of the compounds of formula I will be pharmaceutically acceptable salts. Other salts may, however, be useful in the preparation of the compounds according to the invention or of their
20 pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds of this invention include acid addition salts which may, for example, be formed by mixing a solution of the compound according to the invention with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid, methanesulphonic acid,
25 fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, oxalic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid.

The present invention includes within its scope prodrugs of the compounds of formula I above. In general, such prodrugs will be functional derivatives of the compounds of formula I which are readily
30 convertible *in vivo* into the required compound of formula I. Conventional procedures for the selection and preparation of suitable prodrug

derivatives are described, for example, in *Design of Prodrugs*, ed. H. Bundgaard, Elsevier, 1985.

The compounds according to the invention have at least one asymmetric centre, and they may accordingly exist as enantiomers.

- 5 Where the compounds according to the invention possess two or more asymmetric centres, they may additionally exist as diastereoisomers. It is to be understood that all such isomers and mixtures thereof in any proportion are encompassed within the scope of the present invention.

In the compounds of formula I above, the moiety R¹ suitably
10 represents hydrogen, fluoro, trifluoromethyl, methoxy or a group of formula (a) as defined above. Particular values of R¹ include hydrogen, fluoro and trifluoromethyl, especially hydrogen or fluoro.

Suitably, R² and R³ independently represent hydrogen, fluoro, trifluoromethyl or methoxy, in particular hydrogen or fluoro. Suitably,
15 one or both of R² and R³ represents hydrogen.

Suitably, R⁴ represents methyl, methoxy, isopropoxy or methoxyethoxy.

Specific compounds within the scope of the present invention include:

- 20 4-(1-phenylethyl)-1-[3-(5-(1,2,4-triazol-4-yl)-1*H*-indol-3-yl)propyl]-1,2,5,6-tetrahydropyridine;
4-(α -isopropoxy)phenylmethyl-1-[3-(5-(1,2,4-triazol-4-yl)-1*H*-indol-3-yl)propyl]-1,2,5,6-tetrahydropyridine;
4-(α -methoxy)phenylmethyl-1-[3-(5-(1,2,4-triazol-4-yl)-1*H*-indol-3-yl)propyl]-1,2,5,6-tetrahydropyridine;
25 4-[α -(2-methoxyethyl)oxy]phenylmethyl-1-[3-(5-(1,2,4-triazol-4-yl)-1*H*-indol-3-yl)propyl]-1,2,5,6-tetrahydropyridine;
and salts and prodrugs thereof.


- The invention also provides pharmaceutical compositions
30 comprising one or more compounds of this invention in association with a pharmaceutically acceptable carrier. Preferably these compositions are in

unit dosage forms such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, auto-injector devices or suppositories; for oral, parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. Typical unit dosage forms contain from 1 to 100 mg, for example 1, 2, 5, 10, 25, 50 or 100 mg, of the active ingredient. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.


The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.

In the treatment of migraine, a suitable dosage level is about 0.01 to 250 mg/kg per day, preferably about 0.05 to 100 mg/kg per day, and especially about 0.05 to 5 mg/kg per day. The compounds may be administered on a regimen of 1 to 4 times per day.

The compounds according to the invention may be prepared by a process which comprises reacting a compound of formula II with a compound of formula III:



(II)



(III)


wherein R^1 , R^2 , R^3 and R^4 are as defined above, and L^1 represents a suitable leaving group.

The leaving group L^1 is suitably an alkylsulphonyloxy or arylsulphonyloxy group, e.g. methanesulphonyloxy (mesyloxy) or *p*-toluenesulphonyloxy (tosyloxy).

Where L^1 represents an alkylsulphonyloxy or arylsulphonyloxy group, the reaction between compounds II and III is conveniently carried

out in a suitable solvent such as 1,2-dimethoxyethane or isopropyl alcohol, typically in the presence of a base such as sodium carbonate or potassium carbonate, optionally with the addition of sodium iodide.


In one representative approach, the compounds of formula II wherein L¹ represents a mesyloxy or tosyloxy group may be prepared by the sequence of steps illustrated in the following reaction scheme (cf. Larock and Yum, *J. Am. Chem. Soc.*, 1991, 113, 6689):



wherein L² represents mesyloxy or tosyloxy, and TMS is an abbreviation for trimethylsilyl.

In Step 1 of the reaction scheme, the aniline derivative IV is treated with iodine monochloride, advantageously in methanol in the presence of a base such as calcium carbonate, in order to introduce an iodine atom *ortho* to the amine moiety. Step 2 involves a palladium-mediated coupling reaction with the protected acetylene derivative TMS-C≡C-(CH₂)₃-OH, typically using palladium acetate and triphenylphosphine in the presence of lithium chloride and sodium carbonate, suitably in *N,N*-dimethylformamide at an elevated temperature. This is followed in Step 3 by removal of the TMS moiety, ideally in refluxing methanolic hydrochloric acid; followed in turn by mesylation or tosylation, suitably by using mesyl chloride or tosyl chloride respectively in pyridine.

In another representative approach, the compounds of formula II wherein L¹ represents a mesyloxy or tosyloxy group may be prepared by reacting 3,4-dihydro-2*H*-pyran with the compound of formula V:



(V)

5 followed by mesylation or tosylation of the 3-hydroxypropyl-indole derivative thereby obtained, typically by treatment with mesyl chloride or tosyl chloride under standard conditions.

The reaction with 3,4-dihydro-2H-pyran is suitably brought about by heating the hydrazine derivative V or an acid addition salt thereof, typically the hydrochloride salt, in an inert solvent such as dioxan,
10 advantageously in the presence of a mineral acid such as hydrochloric acid or a Lewis acid such as zinc chloride, at the reflux temperature of the solvent.

The hydrazine derivative of formula V above can be prepared by the method described in WO 94/03446, as also can the aniline derivative of
15 formula IV.

Where they are not commercially available, the starting materials of formula III may be prepared by methods analogous to those described in the accompanying Examples, or by standard procedures well known from the art.

20 Where the above-described processes for the preparation of the compounds according to the invention give rise to mixtures of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The novel compounds may be prepared in racemic form, or individual enantiomers may be prepared
25 either by enantiospecific synthesis or by resolution. The novel compounds may, for example, be resolved into their component enantiomers by standard techniques such as preparative HPLC, or the formation of

diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-*p*-toluoyl-d-tartaric acid and/or (+)-di-*p*-toluoyl-l-tartaric acid, followed by fractional crystallization and regeneration of the free base. The novel compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary.

During any of the above synthetic sequences it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in *Protective Groups in Organic Chemistry*, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

The following Examples illustrate the preparation of compounds according to the invention.

The compounds in accordance with the present invention potently and selectively bind to the 5-HT_{1D α} receptor subtype, inhibit forskolin-stimulated adenylyl cyclase activity, and stimulate [³⁵S]-GTP γ S binding to membranes from clonal cell lines expressing human cloned receptors.

5-HT_{1D α} /5-HT_{1D β} Radioligand Binding

Chinese hamster ovary (CHO) clonal cell lines expressing the human 5-HT_{1D α} and 5-HT_{1D β} receptors were harvested in PBS and homogenised in ice cold 50 mM Tris-HCl (pH 7.7 at room temperature) with a Kinematica polytron and centrifuged at 48,000g at 4°C for 11 min. The pellet was then resuspended in 50 mM Tris-HCl followed by a 10 min incubation at 37°C. Finally the tissue was recentrifuged at 48,000g, 4°C for 11 min and the pellet resuspended, in assay buffer (composition in mM: Tris-HCl 50, pargyline 0.01, CaCl₂ 4; ascorbate 0.1%; pH 7.7 at room

temperature) to give the required volume immediately prior to use (0.2 mg protein/ml). Incubations were carried out for 30 min at 37°C in the presence of 0.02-150 nM [³H]-5-HT for saturation studies or 2-5 nM [³H]-5-HT for displacement studies. The final assay volume was 1 ml. 5-HT (10
5 μM) was used to define non-specific binding. The reaction was initiated by the addition of membrane and was terminated by rapid filtration through Whatman GF/B filters (presoaked in 0.3% PEI/ 0.5% Triton X) followed by 2 x 4 ml washings with 50 mM Tris-HCl. The radioactive filters were then counted on a LKB beta or a Wallac beta plate counter. Binding
10 parameters were determined by non-linear, least squares regression analysis using an iterative curve fitting routine, from which IC₅₀ (the molar concentration of compound necessary to inhibit binding by 50%) values could be calculated for each test compound. The IC₅₀ values for binding to the 5-HT_{1Dα} receptor subtype obtained for the compounds of the
15 accompanying Examples were below 50 nM in each case. Furthermore, the compounds of the accompanying Examples were all found to possess a selective affinity for the 5-HT_{1Dα} receptor subtype of at least 10-fold relative to the 5-HT_{1Dβ} subtype.

20 5-HT_{1Dα}/5-HT_{1Dβ} Adenylyl Cyclase Assay

Studies were performed essentially as described in *J. Pharmacol. Exp. Ther.*, 1986, 238, 248. CHO clonal cell lines expressing the human cloned 5-HT_{1Dα} and 5-HT_{1Dβ} receptors were harvested in PBS and
25 homogenised, using a motor driven teflon/glass homogeniser, in ice cold Tris HCl-EGTA buffer (composition in mM: Tris HCl 10, EGTA 1, pH 8.0 at room temperature) and incubated on ice for 30-60 min. The tissue was then centrifuged at 20,000g for 20 min at 4°C, the supernatant discarded and the pellet resuspended in Tris HCl-EDTA buffer (composition in mM:
30 Tris HCl 50, EDTA 5, pH 7.6 at room temperature) just prior to assay. The adenylyl cyclase activity was determined by measuring the conversion

of α -[^{33}P]-ATP to [^{33}P]-cyclic AMP. A 10 μl aliquot of the membrane suspension was incubated, for 10-15 min, in a final volume of 50 μl , at 30°C, with or without forskolin (10 μM), in the presence or absence of test compound. The incubation buffer consisted of 50 mM Tris HCl (pH 7.6 at room temperature), 100 mM NaCl, 30 μM GTP, 50 μM cyclic AMP, 1 mM dithiothreitol, 1 mM ATP, 5 mM MgCl_2 , 1 mM EGTA, 1 mM 3-isobutyl-1-methylxanthine, 3.5 mM creatinine phosphate, 0.2 mg/ml creatine phosphokinase, 0.5-1 μCi α -[^{33}P]-ATP and 1 nCi [^3H]-cyclic AMP. The incubation was initiated by the addition of membrane, following a 5 min preincubation at 30°C, and was terminated by the addition of 100 μl SDS (composition in mM: sodium lauryl sulphate 2%, ATP 45, cyclic AMP 1.3, pH 7.5 at room temperature). The ATP and cyclic AMP were separated on a double column chromatography system (*Anal. Biochem.*, 1974, 58, 541). Functional parameters were determined using a least squares curve fitting programme ALLFIT (*Am. J. Physiol.*, 1978, 235, E97) from which E_{max} (maximal effect) and EC_{50} (the molar concentration of compound necessary to inhibit the maximal effect by 50%) values were obtained for each test compound. Of those compounds which were tested in this assay, the EC_{50} values for the 5-HT $_{1D\alpha}$ receptor obtained for the compounds of the accompanying Examples were below 500 nM in each case. Moreover, the compounds of the accompanying Examples which were tested were all found to possess at least a 10-fold selectivity for the 5-HT $_{1D\alpha}$ receptor subtype relative to the 5-HT $_{1D\beta}$ subtype.

25 5-HT $_{1D\alpha}$ /5-HT $_{1D\beta}$ GTP γ S Binding

Studies were performed essentially as described in *Br. J. Pharmacol.*, 1993, 109, 1120. CHO clonal cell lines expressing the human cloned 5-HT $_{1D\alpha}$ and 5-HT $_{1D\beta}$ receptors were harvested in PBS and homogenised using a Kinematica polytron in ice cold 20 mM HEPES containing 10 mM EDTA, pH 7.4 at room temperature. The membranes

were then centrifuged at 40,000g, 4°C for 15 min. The pellet was then resuspended in ice cold 20 mM HEPES containing 0.1 mM EDTA, pH 7.4 at room temperature and recentrifuged at 40,000g, 4°C for 15-25 minutes. The membranes were then resuspended in assay buffer (composition in

5 mM: HEPES 20, NaCl 100, MgCl₂ 10, pargyline 0.01; ascorbate 0.1%; pH 7.4 at room temperature) at a concentration of 40 µg protein/ml for the 5-HT_{1D α} receptor transfected cells and 40-50 µg protein/ml for the 5-HT_{1D β} receptor transfected cells. The membrane suspension was then incubated, in a volume of 1 ml, with GDP (100 µM for 5-HT_{1D α} receptor transfected

10 cells, 30 µM for the 5-HT_{1D β} receptor transfected cells) and test compound at 30°C for 20 min and then transferred to ice for a further 15 min. [35S]-GTP γ S was then added at a final concentration of 100 pM and the samples incubated for 30 min at 30°C. The reaction was initiated by the addition of membrane and was terminated by rapid filtration through

15 Whatman GF/B filters and washed with 5 ml water. The radioactive filters were then counted on a LKB beta counter. Functional parameters were determined by a non-linear, least squares regression analysis using an iterative curve fitting routine, from which E_{max} (maximal effect) and EC₅₀ (the molar concentration of compound necessary to inhibit the

20 maximal effect by 50%) values were obtained for each test compound. Of those compounds which were tested in this assay, the EC₅₀ values for the 5-HT_{1D α} receptor obtained for the compounds of the accompanying Examples were below 500 nM in each case. Moreover, the compounds of the accompanying Examples which were tested were all found to possess

25 at least a 10-fold selectivity for the 5-HT_{1D α} receptor subtype relative to the 5-HT_{1D β} subtype.

EXAMPLE 14-(1-Phenylethyl)-1-(3-[5-(1,2,4-triazol-4-yl)-1H-indol-3-yl]propyl)-1,2,5,6-tetrahydropyridine Hydrogen Oxalate

5

a) 4-(1-Phenylethenyl)pyridine

Methyltriphenylphosphonium iodide (12.13g, 30mmol) and potassium t-butoxide (3.37g, 30mmol) in anhydrous tetrahydrofuran (110ml) was stirred and heated at reflux under a nitrogen atmosphere for 1.5 hours. The mixture was cooled to 2°C then treated with a solution of 4-benzoylpyridine (5.0g, 27.3mmol) in anhydrous tetrahydrofuran (30ml) dropwise. The reaction mixture was stirred at 0°C for 1 hour then at room temperature for 18 hours. The mixture was treated with saturated ammonium chloride solution (75ml), the organic layer was collected and the aqueous re-extracted with ethyl acetate (2x100ml). The combined organics were dried (sodium sulphate), evaporated, then the residue was purified by column chromatography on silica using 25% ethyl acetate in petroleum ether (60-80). The title compound was obtained as a gum (3.77g, 76%), δ (360MHz, CDCl₃) 5.60 (2H, d, J=3Hz), 7.24-7.40 (7H, m), 8.60 (2H, d, J=5Hz).

b) 4-(1-Phenylethyl)pyridine

The foregoing olefin (3.75g, 20.7mmol) was hydrogenated at 50 psi in ethanol (50ml) over 10% palladium on carbon. The mixture was filtered and evaporated to give the title compound as a gum (3.28g, 87%), δ (360MHz, CDCl₃) 1.64 (3H, d, J=7Hz), 4.11 (1H, q, J=7Hz), 7.12-7.33 (7H, m), 8.48 (2H, d, J=5Hz).

c) 1-Benzyl-4-(1-phenylethyl)pyridinium bromide

A solution of the foregoing pyridine (3.25g, 17.8mmol) in acetone (25ml) was treated with benzyl bromide (2.21ml, 18.6mmol) and stirred at

room temperature for 18 hours. Diethyl ether (25ml) was added, the solvent decanted and the precipitate triturated with diethyl ether. The precipitate was collected and dried (6.26g, 99%), δ (360MHz, DMSO- d_6) 1.67 (3H, d, $J=7$ Hz), 4.53 (1H, q, $J=7$ Hz), 5.80 (2H, s), 7.25-7.55 (10H, m),
5 8.09 (2H, d, $J=7$ Hz), 9.10 (2H, d, $J=7$ Hz).

d) 1-Benzyl-4-(1-phenylethyl)-1,2,5,6-tetrahydropyridine

The foregoing quaternary salt (6.20g, 17.5mmol) in ethanol (30ml) and water (5ml) at 0°C was treated with sodium borohydride (671mg,
10 17.8mmol). The mixture was allowed to warm to room temperature over 16 hours then treated with a further quantity of sodium borohydride (500mg, 13.2mmol). After a further 16 hours the ethanol was evaporated, the residue diluted with water and extracted with ethyl acetate (4x50ml). The combined organics were dried (sodium sulphate) then evaporated and
15 the residue purified by column chromatography on silica using 25% ethyl acetate in hexane. The title compound was obtained as an oil (3.52g, 72%), δ (360MHz, $CDCl_3$) 1.35 (3H, d, $J=7$ Hz), 1.94-1.98 (2H, m), 2.43-2.55 (2H, m), 3.02-3.10 (2H, m), 3.32 (1H, q, $J=7$ Hz), 3.59 (2H, s), 5.52 (1H, d, $J=2$ Hz), 7.15-7.35 (10H, m).

20

e) 4-(1-Phenylethyl)-1,2,5,6-tetrahydropyridine

A stirred, cooled (0°C) solution of the foregoing amine (2.g, 7.2mmol) in anhydrous dichloromethane (30ml) was treated with α -chloroethyl chloroformate (0.94ml, 8.7mmol). The reaction mixture was allowed to
25 warm to room temperature over 16 hours, then evaporated, dissolved in methanol (30ml) and heated at reflux for 4 hours. The solvent was evaporated, the free base liberated and purified by column chromatography on silica using dichloromethane/methanol/ammonia (9:10:0.1). The title product was obtained as a viscous gum (1.04, 77%), δ
30 (250MHz, $CDCl_3$) 1.36 (3H, d, $J=7$ Hz), 1.83-1.88 (2H, m), 2.82-2.93 (2H,

m), 3.30 (1H, q, J=7Hz), 3.37-3.42 (2H, m), 5.59 (1H, d, J=2Hz), 7.15-7.32 (5H, m). MS, ES⁺, m/z=188 for (M+H)⁺.

f) 4-(1-Phenylethyl)-1-(3-[5-(1,2,4-triazol-4-yl)-1H-indol-3-yl]propyl)-

5 1,2,5,6-tetrahydropyridine Hydrogen Oxalate

A suspension of 3-(5-[1,2,4-triazol-4-yl]-1H-indol-3-yl)propan-1-ol (220mg, 0.91mmol) in anhydrous tetrahydrofuran (30ml), under an atmosphere of nitrogen, was treated with triethylamine (0.25ml, 1.82mmol) and methanesulphonyl chloride (0.14ml, 1.82mmol). The reaction mixture was stirred for 1.5 hours, filtered, evaporated then redissolved in dichloromethane (50ml) and washed with water (2x30ml). The combined organics were dried (sodium sulphate), evaporated and the crude mesylate was suspended in propan-2-ol (50ml) then treated with potassium carbonate (500mg, 3.6mmol) and 4-(1-phenylethyl-1,2,5,6-tetrahydropyridine (482mg, 2.5mmol). The reaction mixture was stirred whilst heating at reflux for 66 hours, then cooled and evaporated. The residue was partitioned between water (50ml) and dichloromethane (50ml). The aqueous was separated then re-extracted with dichloromethane (3x50ml). The combined organics were dried (sodium sulphate), evaporated and the residue purified by chromatography on silica using dichloromethane/methanol/ammonia (95:4:1) to give the title compound free base (115mg, 31%). The hydrogen oxalate salt had mp >120°C (dec.). MS, ES⁺, m/z=412 for (M+H)⁺, δ (360MHz, DMSO-d₆) 1.31 (3H, d, J=7Hz), 2.02-2.20 (4H, m), 2.78 (2H, t, J=7Hz), 3.05-3.30 (4H, m), 3.45 (1H, q, J=7Hz), 3.60-3.80 (2H, m), 5.56 (1H, s), 7.17-7.25 (3H, m), 7.27-7.35 (4H, m), 7.51 (1H, d, J=9Hz), 7.80 (1H, d, J=2Hz), 9.01 (2H, s), 11.20 (1H, s). (Found: C, 63.89; H, 6.23; N, 12.24. C₂₆H₂₉N₅. 1.45C₂H₂O₄. 0.33CH₃CH₂OH requires C, 63.71; H, 6.13; N, 12.57%).

EXAMPLE 2

4-(α -Isopropoxy)phenylmethyl-1-(3-[5-(1,2,4-triazol-4-yl)-1H-indol-3-yl]propyl)-1,2,5,6-tetrahydropyridine Hydrogen Oxalate

a) α -Phenyl- α -(4-pyridyl)methanol

A 1M solution of phenylmagnesium bromide in tetrahydrofuran (100ml, 100mmol) was treated with 4-pyridylcarboxaldehyde (10g, 93.5mmol) in anhydrous tetrahydrofuran (50ml) under a nitrogen atmosphere. The mixture was heated at reflux with stirring for 4 hours, cooled to room temperature then treated with saturated ammonium chloride solution followed by dichloromethane then water. The organic layer was separated and the aqueous re-extracted with dichloromethane. The combined organics were dried (sodium sulphate) then evaporated then the residue triturated with hexane/ethyl acetate (3:1) to give the required product (13.5g, 73%). MS, ES⁺, m/z=186 for (M+H)⁺.

b) α -(1-Benzyl-1,2,5,6-tetrahydropyridin-4-yl)- α -phenylmethanol

The foregoing pyridine (10g, 54mmol) in acetone (50ml) was treated with benzyl bromide (6.4ml, 54mmol) and the reaction mixture was stirred at room temperature for 18 hours. The oil produced was decanted and triturated with acetone to give the quaternary salt (19g, 98%). This quaternary salt (12.0g, 33.7mmol) in ethanol (500ml) was treated portionwise with sodium borohydride (1.5g, 40mmol). The reaction mixture was stirred at room temperature for 18 hours. The reaction mixture was treated with water, then the majority of the solvent evaporated and the residue extracted into dichloromethane. The organic layer was separated and the aqueous re-extracted with dichloromethane. The combined organics were dried (sodium sulphate) then evaporated and the residue purified by column chromatography on silica using

dichloromethane/methanol (9:1) to give the title tetrahydropyridine (7.2g, 77%), δ (250MHz, CDCl_3) 1.97-2.05 (2H, m), 2.52 (2H, dd, $J_1=J_2=6\text{Hz}$), 3.01-3.06 (2H, m), 3.57 (2H, s), 5.10 (1H, s), 5.75-5.79 (1H, m), 7.20-7.37 (10H, m).

5

c) O-Isopropyl- α -phenyl- α -(1,2,5,6-tetrahydropyridin-4-yl)methanol

The foregoing tetrahydropyridine (2.0g, 7.1mmol) in dichloromethane (100ml), at 0°C , was treated with α -chloroethyl chloroformate (1.7ml, 16mmol). The solution was stirred, whilst warming to room temperature, for 18 hours. The solvent was evaporated then the residue treated with propan-2-ol (100ml) and heated at reflux for 1 hour. The solvent was evaporated and the residue partitioned between saturated aqueous potassium carbonate solution and dichloromethane. The organic layer was separated and the aqueous re-extracted with dichloromethane. The combined organics were dried (sodium sulphate), evaporated then the residue was purified by column chromatography on neutral alumina using dichloromethane/methanol (9:1). The required tetrahydropyridine was obtained as a gum (620mg, 38%), δ (250MHz, CDCl_3) 1.14 and 1.20 (each 3H, each d, $J=6\text{Hz}$), 1.90-1.98 (2H, m), 2.92 (2H, dd, $J_1=J_2=6\text{Hz}$), 3.39 (2H, d, $J=3\text{Hz}$), 3.63 (1H, heptet, $J=6\text{Hz}$), 4.78 (1H, s), 5.70-5.73 (1H, m), 7.23-7.34 (5H, m), MS, ES^+ , $m/z=232$ for $(\text{M}+\text{H})^+$.

20

d) 4-(α -Isopropoxy)phenylmethyl-1-(3-[5-(1,2,4-triazol-4-yl)-1H-indol-3-yl]propyl)-1,2,5,6-tetrahydropyridine Hydrogen Oxalate

The title compound free base was obtained (80mg, 21%) from O-isopropyl- α -phenyl- α -(1,2,5,6-tetrahydropyridin-4-yl)methanol and the mesylate obtained from 3-(5-[1,2,4-triazol-4-yl]-1H-indol-3-yl)propan-1-ol as described in Example 1. The hydrogen oxalate salt has mp $>122^\circ\text{C}$ (dec.). MS, ES^+ , $m/z=456$ for $(\text{M}+\text{H})^+$, δ (360MHz, $\text{DMSO}-d_6$) 1.07 and 1.13 (each 3H, each d, $J=6\text{Hz}$), 2.00-2.20 (4H, m), 2.76 (2H, t, $J=7\text{Hz}$), 3.04-3.35 (4H, m), 3.54 (1H, heptet, $J=6\text{Hz}$), 3.73 (2H, br s), 4.96 (1H, s), 5.77 (1H, s),

30

7.26-7.36 (7H, m), 7.50 (1H, d, J=8Hz), 7.79 (1H, d, J=2Hz), 9.01 (2H, s), 11.19 (1H, s). (Found C, 60.80; H, 6.04; N, 11.44. $C_{28}H_{33}N_5O \cdot 1.9C_2H_2O_4$ requires C, 60.95; H, 5.92; N, 11.18%).

5

EXAMPLE 3

4-(α -Methoxy)phenylmethyl-1-(3-[5-(1,2,4-triazol-4-yl)-1H-indol-3-yl]propyl)-1,2,5,6-tetrahydropyridine Hydrogen Oxalate

- 10 a) **O-Methyl- α -phenyl- α -(1,2,5,6-tetrahydropyridin-4-yl)methanol**
 α -(1-Benzyl-1,2,5,6-tetrahydropyridin-4-yl)- α -phenylmethanol (7.0g, 25mmol) in dichloromethane (200ml) was cooled to -5°C then treated with α -chloroethyl chloroformate (6ml, 55mmol). The reaction mixture was stirred whilst warming to room temperature over 18 hours. The solvent
15 was evaporated then the residue was dissolved in methanol (200ml) and heated at reflux for 3 hours. The solvent was evaporated, the free base was liberated and purified by column chromatography on neutral alumina using dichloromethane/methanol (9:1). The title compound was obtained as a gum (2.0g, 39%). MS, ES⁺, m/z=204 for (M+H)⁺, δ (250MHz, CDCl₃)
20 1.80-1.96 (2H, m), 2.86-2.95 (2H, m), 3.32 (3H, s), 3.37-3.41 (2H, m), 4.54 (2H, s), 5.78-5.84 (1H, m), 7.21-7.40 (5H, m).

b) **4-(α -Methoxy)phenylmethyl-1-(3-[5-(1,2,4-triazol-4-yl)-1H-indol-3-yl]propyl)-1,2,5,6-tetrahydropyridine Hydrogen Oxalate**

- 25 The title compound free base was obtained (150mg, 30%) from the foregoing tetrahydropyridine and the mesylate obtained from 3-(5-[1,2,4-triazol-4-yl]-1H-indol-3-yl)propan-1-ol as described in Example 1. The hydrogen oxalate salt had mp >90°C. MS, ES⁺, m/z=428 for (M+H)⁺. (Found: C, 59.21; H, 5.76; N, 11.68. $C_{26}H_{29}N_5O \cdot 2C_2H_2O_4$ requires C, 59.30;
30 H, 5.47; N, 11.52%).

EXAMPLE 4

5 4-(α -[2-Methoxyethyl]oxy)phenylmethyl-1-(3-[5-(1,2,4-triazol-4-yl)-1H-indol-3-yl]propyl)-1,2,5,6-tetrahydropyridine Hydrogen Oxalate

a) O-(2-Methoxyethyl)- α -phenyl- α -(1,2,5,6-tetrahydropyridin-4-yl)methanol

10 The title compound was obtained (250mg, 29%) from α -(1-benzyl-1,2,5,6-tetrahydropyridin-4-yl)- α -phenylmethanol using 2-methoxyethanol instead of propan-2-ol, as described in Example 2. MS, ES⁺, m/z=248 for (M+H)⁺, δ (250MHz, CDCl₃) 1.73-1.77 (2H, m), 2.90 (2H, dd, J=6Hz), 3.38-3.42 (5H, m), 3.50-3.62 (4H, m), 4.70 (1H, s), 5.78-5.80 (1H, m), 7.24-7.34 (5H, m).


15

b) 4-(α -[2-Methoxyethyl]oxy)phenylmethyl-1-(3-[5-(1,2,4-triazol-4-yl)-1H-indol-3-yl]propyl)-1,2,5,6-tetrahydropyridine Hydrogen Oxalate

20 The title compound free base was obtained (100mg, 26%) from the foregoing tetrahydropyridine and the mesylate obtained from 3-(5-[1,2,4-triazol-4-yl]-1H-indol-3-yl)propan-1-ol as described in Example 1. The hydrogen oxalate salt had mp >109°C (dec.). MS, ES⁺, m/z=472 for (M+H)⁺. (Found: C, 61.00; H, 6.91; N, 11.50. C₂₈H₃₃N₅O₂. C₂H₂O₄. 1.5H₂O requires C, 61.21; H, 6.51; 11.90%).

CLAIMS:


1. A compound of formula I, or a salt or prodrug thereof:



(I)

wherein

R¹ represents hydrogen, halogen, trifluoromethyl, C₁₋₆ alkoxy or a group of formula (a):



(a)

R² and R³ independently represent hydrogen, halogen, trifluoromethyl or C₁₋₆ alkoxy; and

R⁴ represents C₁₋₆ alkyl, C₁₋₆ alkoxy or C₁₋₆ alkoxy(C₁₋₆)alkoxy.

2. A compound as claimed in claim 1 wherein R¹ represents hydrogen or fluoro.

3. A compound as claimed in claim 1 or claim 2 wherein R² and R³ independently represent hydrogen or fluoro.

4. A compound as claimed in any one of the preceding claims wherein R¹, R² and R³ each represents hydrogen.

5. A compound as claimed in any one of the preceding claims
5 wherein R⁴ represents methyl, methoxy, isopropoxy or methoxyethoxy.

6. A compound selected from:

4-(1-phenylethyl)-1-[3-(5-(1,2,4-triazol-4-yl)-1*H*-indol-3-yl)propyl]-1,2,5,6-tetrahydropyridine;

10 4-(α -isopropoxy)phenylmethyl-1-[3-(5-(1,2,4-triazol-4-yl)-1*H*-indol-3-yl)propyl]-1,2,5,6-tetrahydropyridine;

4-(α -methoxy)phenylmethyl-1-[3-(5-(1,2,4-triazol-4-yl)-1*H*-indol-3-yl)propyl]-1,2,5,6-tetrahydropyridine;

15 4-[α -(2-methoxyethyl)oxy]phenylmethyl-1-[3-(5-(1,2,4-triazol-4-yl)-1*H*-indol-3-yl)propyl]-1,2,5,6-tetrahydropyridine;
and salts and prodrugs thereof

7. A pharmaceutical composition comprising a compound of formula I as defined in claim 1 or a pharmaceutically acceptable salt
20 thereof or a prodrug thereof in association with a pharmaceutically acceptable carrier.


8. A compound as claimed in any one of claims 1 to 6 for use in therapy.

25


9. The use of a compound as claimed in any one of claims 1 to 6 for the manufacture of a medicament for the treatment and/or prevention of clinical conditions for which an agonist of 5-HT_{1D} receptors selective for the 5-HT_{1D α} subtype thereof is indicated.

30

10. A process for the preparation of a compound as claimed in claim 1, which comprises reacting a compound of formula II with a compound of formula III:



(II)



(III)

wherein R¹, R², R³ and R⁴ are as defined in claim 1, and L¹ represents a suitable leaving group.

- 10 11. A method for the treatment and/or prevention of clinical conditions for which an agonist of 5-HT_{1D} receptors selective for the 5-HT_{1D_α} subtype thereof is indicated, which method comprises administering to a patient in need of such treatment an effective amount of a compound of formula I as defined in claim 1, or a pharmaceutically acceptable salt thereof or a prodrug thereof.
- 15

INTERNATIONAL SEARCH REPORT

Intern. Application No
PCT/GB 96/02684

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D401/14 A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94 02477 A (MERCK SHARP & DOHME LTD.) 3 February 1994 cited in the application see claims ---	1-11
P,A	WO 95 32196 A (MERCK SHARP & DOHME LTD.) 30 November 1995 see claims ---	1-11
P,A	WO 96 16056 A (MERCK SHARP & DOHME LTD.) 30 May 1996 see claims -----	1-11

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *A* document member of the same patent family

Date of the actual completion of the international search

14 January 1997

Date of mailing of the international search report

28.01.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Chouly, J

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB 96/02684

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 11 is directed to a method of treatment of (diagnostic method practised on) the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Interns Application No
PCT/GB 96/02684

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9402477	03-02-94	AU-B- 672802	17-10-96
		AU-A- 4578593	14-02-94
		CA-A- 2138649	03-02-94
		EP-A- 0651749	10-05-95
		JP-T- 7509452	19-10-95
		US-A- 5567726	22-10-96
-----	-----	-----	-----
WO-A-9532196	30-11-95	AU-A- 2529695	18-12-95
-----	-----	-----	-----
WO-A-9616056	30-05-96	AU-A- 3875095	17-06-96
-----	-----	-----	-----